

Final Episode Report

Christiaan Barnard Memorial Hospital
PathCare Reference Laboratory
PathCare Park
Room 1619, 16th Floor
Cnr DF Malan and Rua Bartholomeu Dias Plain
Neels Bothma Street
Tel : +27 21 596 3400
Tel: 021 424 5102



Practice No:0774383

Report to:
DALING JAN-MARTEN

Referred by: DR JILL HARRIS
Copies to: DR J C H LAPORTA

Requisition No: 654371236

Patient: (File No: 2454217)

Guarantor:

Collection Date: 2025-07-07 09:50

MR JAN-MARTEN DALING

MR J M DALING

Received Date: 2025-07-07 11:17

Patient ID No: 8305145088089

Med Aid: DISCOVERY

Generated On: 2025-07-15 19:01

Age:Sex:DoB: 42y: M: 1983-05-14

Member No: 255751841

Contact No: 0825578133

Contact No: 0825578133

Patient Email: JMDALING@GMAIL.COM

Tests requested: HAEMOGLOBIN + MCV; PLATELET COUNT; INR; C-REACTIVE PROTEIN; VITAMIN D3 (25 OH)

Referral ICD10 C37
code(s):

Ch Biochemistry

0707:BA03221U

Final

Test Name

C-REACTIVE PROTEIN

Result

128.2

Flag

H

Reference Range

0-5.0 mg/L

*** Delta : 14.8 - May 23 2025 4:03PM

En Endocrinology

0707:EA01350U

Final

Test Name

VITAMIN D (25 OH) (ABBOTT)

Result

40

Reference Range

ng/mL

Interpretation of 25-OH vit d level [ng/mL]:

Deficiency: < 12

Partial deficiency: 12 - 19

Optimal level: > 20

Toxicity: > 100

Ref: Munn et al. JCEM.2016;101(2):394

Ha Haematology

0707:HA02525U

Final

Test Name

Result

Flag

Reference Range

RED CELLS

Haemoglobin

10.8

L

12.5 - 16.5 g/dL

MCV

95

#

81 - 95 fl

PLATELETS

Platelet count

197

140 - 420 x10E9/L

Prothrombin Time

11.70

9.9 - 11.8 sec

Control time	10.70	sec
	NOTE: The Prothrombin Time measures the overall efficiency of the extrinsic clotting system. Common causes of a prolonged result include anticoagulation therapy, liver disease, vitamin K deficiency, disseminated intravascular coagulation or congenital factor deficiencies. The Prothrombin Time is used to calculate the INR in patients on warfarin therapy.	
INR	1.10	

NOTE: This is a direct INR. The INR is calculated from a calibration line.

GENERAL GUIDELINES FOR PATIENTS ON WARFARIN THERAPY

The target INR is 2.5 (range 2.0 - 3.0) for most indications (including low-risk patient with bi-leaflet mechanical valves such as the St Jude Medical device in the aortic position) and 3.0 (range 2.5 - 3.5) for other mechanical prosthetic valves.

The interpretation of laboratory test results requires the clinical evaluation to be known and contextualised. Please contact your medical practitioner for any questions related to these results. Your doctor would know whether further consultation with one of our specialist pathologists is necessary.

L=Low *L=Critically Low H=High *H=Critically High #=Delta Checked